Datasheet

Blood coagulation and Anemia • Bone Metabolism • Cardiac Markers • Fertility and Pregnancy Cangliosides • Hormone Markers • Immunology and Serology • Infectious Diseases • Inflammation Kidney Diseases • Metabolic Syndrome • Microbial and Plant Toxins • Miscellaneous • Neuroscience Thyroid Diseases • Tumor Markers • Veterinary

CATALOGUE #: 8T53

PRODUCT NAME: Human cardiac troponin I (cTnI)

Source:	Human heart tissue.
	Blood sample from the tissue donors was tested and found negative for HBsAg, HIV-1 and HIV-2 antibodies, HCV, and syphilis.
Applications:	Tnl is the inhibitory subunit of troponin, the thin filament regulatory complex which confers calcium sensitivity to striated muscle actomyosin ATPase activity.
	Tnl exists in three isoforms, one for slow-twitch skeletal muscle, one for fast twitch skeletal muscle and one for cardiac muscle. The amino acid sequences of two skeletal (skTnl) and one cTnl forms show 40% dissimilarity. Furthermore, the human cTnl has additional amino acid residues on its N-terminal region that are not present on the skeletal form, making this protein a promising analyte for indicating cardiac specificity. cTnl follows similar release pattern as CK-MB within the first 15 hours of myocardial infarction (MI), but remains elevated in serum up to 6 days. This observation and the unique amino sequence of cTnl make it an important diagnostic marker of MI.
	TnI is suitable for use as a standard in immunoassay for early detection of MI, immunogen for antiserum production and tracer for iodination.
Analysis:	Purity > 98 % (SDS-PAGE).
	Immunological activity confirmed by reaction with monoclonal antibody that is specific for the troponin I cardiac isoform in immunoblotting.
	Tnl concentration was determined spectrophotometrically using A (0.1 %, 280 nm, 1 cm) equal to 0.42. This coefficient was calculated from the amino composition of human cTnl (FEBS Lett, 270, 57-61).
Presentation:	Lyophilized from 0.01 M HCI.
	It is recommended to reconstitute this product with Tris/urea buffer (20 mM Tris, pH 7.5, 7 M urea, 5 mM EDTA, 15 mM 2-mercaptoethanol) to a concentration close to 1 mg/ml.
	Solubility is limited at neutral pH and physiological salt concentration.
Storage:	Lyophilized -20°C (-1530 °C allowed)
	Reconstituted -70°C (-6580 °C allowed)
Other information:	Avoid repeated freezing and thawing. It is recommended to aliquot the product after reconstitution.
Material safety note:	This product is sold for research or further manufacturing use only . Standard Laboratory Practices should be followed when handling this material.



SCIENTIFIC EXCELLENCE FOR IVD

HYTEST LTD

Intelligate 1, 6th floor, Joukahaisenkatu 6 • FI-20520 Turku, FINLAND Tel. +358 2 512 0900 • E-mail: hytest@hytest.fi • **HYTEST.FI**