



Myoglobin



Myoglobin is a small heme-containing protein (153 a.a.r., MW (w/o heme) 17053 Da and theoretical pI=7.29), that is responsible for the oxygen deposition in muscle tissues. Only one form of myoglobin is expressed in cardiac and skeletal muscles.

Myoglobin has been used as a marker of myocardial damage for almost six decades (1). It is commonly used in clinical practice as an



CLINICAL UTILITY

- ✓ Myocardial damage
- ✓ Acute myocardial infarction

early marker of AMI (2). However, due to the high concentration of myoglobin in skeletal muscle tissues, even minor skeletal muscle injury increases the myoglobin levels in blood (3). Therefore, myoglobin alone is not considered to be a reliable and sufficient marker in AMI diagnosis. Instead, it should be used together with cTnI or cTnT analysis as part of a multimarker strategy.

Human myoglobin

Myoglobin is purified from human cardiac tissue by several chromatographic steps including gel-filtration and anion-exchange chromatography.

After SDS-PAGE in reducing conditions myoglobin is presented by a single band with apparent molecular mass of 17 kDa (Fig. 1).

In 2001 HyTest's myoglobin preparation was selected by International Federation of Clinical Chemistry as an International Standard Material.




Figure 1. SDS-PAGE of human myoglobin.

M is the molecular weight standard (Pharmacia).

Monoclonal antibodies specific to myoglobin

Hybridoma clones have been derived from hybridization of Sp2/O myeloma cells with spleen cells of Balb/c mice immunized with purified human myoglobin. Different combinations of monoclonal antibodies can be used for the immunoassay development. The best MAb combinations for sandwich immunoassay are:

4E2 - 7C3
7C3 - 4E2

Myoglobin free serum

Myoglobin free serum is prepared from pooled normal human serum by immunoaffinity chromatography. The matrix for affinity sorbent utilizes several monoclonal antibodies with different epitope specificity. Based on ELISA testing the level of myoglobin in myoglobin free serum is below 0.1 ng/ml (Fig. 2).

Myoglobin free serum is used as a matrix for standard and calibrator preparations.




Figure 2. Myoglobin level in normal human serum and in myoglobin free serum detected in sandwich-immunoassay. Column 1: Buffer, column 2: Pooled normal human serum before myoglobin extraction (4 ng/ml of myoglobin) and column 3: Residual signal in myoglobin free serum.

Ordering information

MONOCLONAL ANTIBODIES

Product name	Cat. #	MAb	Subclass	Remarks
Myoglobin	4M23	4E2	IgG1	EIA
		4E2cc	IgG1	<i>In vitro</i> , EIA
		7C3	IgG1	EIA
		1B4	IgG1	EIA

New!

ANTIGEN

Product name	Cat. #	Purity	Source
Myoglobin	8M50	>95%	Human cardiac muscle

DEPLETED SERUM

Product name	Cat. #	Source
Myoglobin free serum	8MFS	Pooled normal human serum

References

- Kiss A and Reinhart W.** (1956) Determination of myoglobin in serum and in urine after heart infarction. Wien Klin. Wochenschr. 68(9):154-155.
- Penttila K. et al.** (2002) Myoglobin, creatine kinase MB isoforms and creatine kinase MB mass in early diagnosis of myocardial infarction in patients with acute chest pain. Clin Biochem. 35(8), 647-653.
- Van Nieuwenhoven FA, et al.** (1995) Discrimination between myocardial and skeletal muscle injury by assessment of the plasma ratio of myoglobin over fatty acid-binding protein. Circulation. 92(10), 2848-2854.