

## DataSheet



## CATALOGUE #: 8FS5

## PRODUCT NAME: Recombinant feline serum amyloid A (SAA)

| Source:               | Expressed in <i>E.coli</i>  |
|-----------------------|---|
| oource.               |   |
| Description:          | Recombinant SAA consists of 121 amino acids and has a predicted molecular mass of 13838 kDa.  |
|                       | Consists of a polypeptide corresponding to 1-111 a.a.r. of feline SAA and a proprietary 10 a.a.r. affinity tag on the N-terminus.   |
| Applications:         | Can be used as a control in immunoassays.   |
| Analysis:             | Purity >95 % (Tricine-SDS-PAGE).  |
|                       | Concentration is determined spectrophotometrically  |
| Purification:         | Affinity and reversed-phase chromatography  |
| Presentation:         | Lyophilized from 0.01 M HCI (pH 2.0)  |
|                       | It is recommended to reconstitute this product with 0.01 M HCI (pH 2.0) to 0.1-1 mg/ml concentration.   |
| Storage:              | Lyophilized -20°C (-1530 °C allowed)  |
|                       | Reconstituted -70°C (-6580 °C allowed)  |
| Other information:    | Avoid repeated freezing and thawing. It is recommended to aliquot the product after reconstitution.   |
|                       | To prepare diluted samples of SAA it is recommended to use buffer solutions containing a detergent.<br>The selection of an appropriate detergent and its concentration is essential for the assay<br>performance. |
|                       | For sandwich immunoassays using monoclonal anti-SAA antibodies SAA19, SAA21 and VSA34 we recommend using 0.01% CHAPS in antigen dilution buffer (e.g. 0.05 M Tris-HCI, pH 8.0, 0.15 M NaCl, 0.01 % CHAPS).        |
|                       | For sandwich immunoassays using monoclonal anti-SAA antibodies VSA25 and VSA31 we recommend using 0.05 % Tween 20 in antigen dilution buffer (e.g. 0.02 M Tris-HCI, pH 7.5, 0.15 M NaCI, 0.05 % Tween 20).        |
| Material safety note: | This product is sold <b>for research or further manufacturing use only</b> . Standard Laboratory Practices should be followed when handling this material.  |